



4. Molecular dynamics

Molecular Simulations

➔ **Molecular dynamics:**
solve equations of motion



➔ **Monte Carlo:** importance
sampling



Molecular Dynamics

4. Molecular Dynamics

4.1. Introduction

4.2. Basics

4.3. Liouville formulation

4.4. Multiple time steps

4. Molecular dynamics

4.2 Basics

“Fundamentals”

Theory:

$$F = m \frac{d^2 r}{dt^2}$$

- Compute the forces on the particles
- Solve the equations of motion
- Sample after some # of timesteps

4. Molecular dynamics

4.3 Some practical details

Molecular Dynamics

Initialization

- Total momentum should be zero (no external forces)
- Temperature rescaling to desired temperature
- Particles start on a lattice

Force calculations

- Periodic boundary conditions
- Order N^2 algorithm,
- Order N : neighbor lists, linked cell
- Truncation and shift of the potential

Integrating the equations of motion

- Velocity Verlet
- Kinetic energy

Molecular Dynamics

Algorithm 3 (A Simple Molecular Dynamics Program)

```
program md
```

```
call init
```

```
t=0
```

```
do while (t.lt.tmax)
```

```
    call force(f,en)
```

```
    call integrate(f,en)
```

```
    t=t+delt
```

```
    call sample
```

```
enddo
```

```
stop
```

```
end
```

simple MD program

initialization

MD loop

determine the forces

integrate equations of motion

sample averages

3. Molecular dynamics: practical details

3.3.1 Initialization

Algorithm 4 (Initialization of a Molecular Dynamics Program)

<pre>subroutine init sumv=0 sumv2=0 do i=1,npart x(i)=lattice_pos(i) v(i)=(ranf()-0.5) sumv=sumv+v(i) sumv2=sumv2+v(i)**2 enddo sumv=sumv/npart sumv2=sumv2/npart fs=sqrt(3*temp/sumv2) do i=1,npart v(i)=(v(i)-sumv)*fs xm(i)=x(i)-v(i)*dt enddo return end</pre>	<p>initialization of MD program</p> <p>place the particles on a lattice give random velocities velocity center of mass kinetic energy</p> <p>velocity center of mass mean-squared velocity scale factor of the velocities set desired kinetic energy and set velocity center of mass to zero position previous time step</p>
--	--


3. Molecular dynamics: practical details

3.3.2 Force calculation

Algorithm 5 (Calculation of the Forces)

subroutine force(f,en)	determine the force and energy
en=0	
do i=1,npart	
f(i)=0	set forces to zero
enddo	
do i=1,npart-1	
do j=i+1,npart	loop over all pairs
xr=x(i)-x(j)	
xr=xr-box*nint(xr/box)	periodic boundary conditions
r2=xr**2	
if (r2.lt.rc2) then	test cutoff
r2i=1/r2	
r6i=r2i**3	
ff=48*r2i*r6i*(r6i-0.5)	Lennard-Jones potential
f(i)=f(i)+ff*xr	update force
f(j)=f(j)-ff*xr	
en=en+4*r6i*(r6i-1)-ecut	update energy
endif	
enddo	
enddo	
return	
end	

Periodic boundary conditions



The Lennard-Jones potential **s**

- The Lennard-Jones potential

$$U^L(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right]$$


- The truncated Lennard-Jones potential

$$U_{TR}^L(r) = \begin{cases} U^L(r) & r \leq r_c \\ 0 & r > r_c \end{cases}$$

- The truncated and shifted Lennard-Jones potential


$$U_{TR-SH}^L(r) = \begin{cases} U^L(r) - U^L(r_c) & r \leq r_c \\ 0 & r > r_c \end{cases}$$

The Lennard-Jones potentials




Saving CPU-time

Cell list



Verlet-list



3. Molecular dynamics: practical details

3.3.3 Equations of motion

Algorithm 6 (Integrating the Equations of Motion)

```
subroutine integrate(f,en)
sumv=0
sumv2=0
do i=1,npart
  xx=2*x(i)-xm(i)+delt**2*f(i)
  vi=(xx-xm(i))/(2*delt)
  sumv=sumv+vi
  sumv2=sumv2+vi**2
  xm(i)=x(i)
  x(i)=xx
enddo
temp=sumv2/(3*npart)
etot=(en+0.5*sumv2)/npart
return
end
```

integrate equations of motion

MD loop

Verlet algorithm (4.2.3)

velocity (4.2.4)

velocity center of mass

total kinetic energy

update positions previous time

update positions current time

instantaneous temperature

total energy per particle

Equations of motion

We can make a Taylor expansion for the positions:

$$r(t + \Delta t) = r(t) + \frac{dr(t)}{dt} \Delta t + \frac{d^2r(t)}{dt^2} \frac{\Delta t^2}{2!} + O(\Delta t^3)$$

The simplest form (Euler):

$$r(t + \Delta t) = r(t) + v(t) \Delta t + O(\Delta t^2)$$

$$v(t + \Delta t) = v(t) + m \frac{df(t)}{dt} \Delta t$$

We can do better!

We can make a Taylor expansion for the positions:

$$r(t + \Delta t) = r(t) + \frac{dr(t)}{dt} \Delta t + \frac{d^2r(t)}{dt^2} \frac{\Delta t^2}{2!} + \frac{d^3r(t)}{dt^3} \frac{\Delta t^3}{3!} + O(\Delta t^4)$$

$$r(t - \Delta t) = r(t) - \frac{dr(t)}{dt} \Delta t + \frac{d^2r(t)}{dt^2} \frac{\Delta t^2}{2!} - \frac{d^3r(t)}{dt^3} \frac{\Delta t^3}{3!} + O(\Delta t^4)$$

When we add the two:

$$r(t + \Delta t) + r(t - \Delta t) = 2r(t) + \frac{d^2r(t)}{dt^2} \Delta t^2 + O(\Delta t^4)$$

Verlet algorithm

$$r(t + \Delta t) = 2r(t) - r(t - \Delta t) + f(t) \frac{\Delta t^2}{m} + O(\Delta t^4)$$

numerically not
ideal

no need for
velocities

Verlet algorithm:
$$r(t + \Delta t) = 2r(t) - r(t - \Delta t) + f(t) \frac{\Delta t^2}{m} + O(\Delta t^4)$$

Velocity Verlet algorithm

$$r(t + \Delta t) = r(t) + v(t) \Delta t + f(t) \frac{\Delta t^2}{2m} + O(\Delta t^4)$$

$$v(t + \Delta t) = v(t) + \frac{\Delta t}{2m} [f(t + \Delta t) + f(t)]$$

to see the equivalence:

$$r(t + 2\Delta t) = r(t + \Delta t) + v(t + \Delta t) \Delta t + f(t + \Delta t) \frac{\Delta t^2}{2m}$$

$$r(t) = r(t + \Delta t) - v(t) \Delta t - f(t) \frac{\Delta t^2}{2m}$$

adding the two

$$r(t + 2\Delta t) = 2r(t + \Delta t) - r(t) + [v(t + \Delta t) - v(t)] \Delta t + [f(t + \Delta t) - f(t)] \frac{\Delta t^2}{2m}$$

with
$$v(t + \Delta t) = v(t) + \frac{\Delta t}{2m} [f(t + \Delta t) + f(t)]$$

$$r(t + 2\Delta t) = 2r(t + \Delta t) - r(t) + f(t + \Delta t) \frac{\Delta t^2}{m}$$


Lyapunov instability

MD: reference trajectory
with initial condition:

$$(r_1(0), \dots, r_N(0), p_1(0), \dots, p_N(0))$$

MD: compare: $(r_1(0), \dots, r_N(0), p_1(0), \dots, p_i(0) + \varepsilon, p_j(0) - \varepsilon, \dots, p_N(0))$

$$\varepsilon = 10^{-10}$$



4. Molecular dynamics:

4.4 Liouville Formulation

Liouville formulation

the dot above, \dot{f} ,
implies time derivative

Let us consider a function that f which depends on the positions and momenta of the particles:

$$f(p^N, r^N)$$

We can “solve” how f depends on time:

$$\dot{f} = \left(\frac{\partial f}{\partial r} \right) \dot{r} + \left(\frac{\partial f}{\partial p} \right) \dot{p}$$

Define the Liouville operator:

$$iL \equiv \dot{r} \left(\frac{\partial}{\partial r} \right) + \dot{p} \left(\frac{\partial}{\partial p} \right)$$

the time dependence follows from:

$$\frac{df}{dt} = iL f$$

with solution:

beware: the solution is
equally useless as the
differential equation

$$f = e^{iLt} f(0)$$

In an ideal world it would be less useless:

$$iL \equiv \dot{r} \left(\frac{\partial}{\partial r} \right) + \dot{p} \left(\frac{\partial}{\partial p} \right)$$

Let us look at half the equation
which has as solution:

$$iL_r \equiv \left(\frac{\partial}{\partial r} \right) \dot{r}$$

$$f = e^{iL_r t} f(0)$$

Taylor expansion:

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$e^{iL_r t} f(0) = \left[1 + iL_r t + \frac{1}{2} (iL_r t)^2 + \frac{1}{3!} (iL_r t)^3 + \dots \right] f(0)$$

$$e^{iL_r t} f(0) = \left[1 + \dot{r}(0)t \left(\frac{\partial}{\partial r} \right) + \frac{1}{2} (\dot{r}(0)t)^2 \left(\frac{\partial}{\partial r} \right)^2 + \dots \right] f(0)$$

the operator iL_r
gives a shift of
the positions

$$f(0 + \dot{r}(0)t) = f(0) + \dot{r}(0)t \left(\frac{\partial f(0)}{\partial r} \right) + \frac{1}{2} (\dot{r}(0)t)^2 \left(\frac{\partial^2 f(0)}{\partial r^2} \right) + \dots$$

Hence:

$$e^{iL_r t} f(0) = f(0 + \dot{r}(0)t)$$

The operation iL_r gives a shift of the positions

$$iL \equiv \dot{r} \left(\frac{\partial}{\partial r} \right) + \dot{p} \left(\frac{\partial}{\partial p} \right)$$

Similarly for the operator iL_p

$$iL_p \equiv \left(\frac{\partial}{\partial p} \right) \dot{p}$$

which has as solution:

$$f = e^{iL_p t} f(0)$$

Taylor expansion:

$$e^{iL_p t} f(0) = \left[1 + iL_p t + \frac{1}{2} (iL_p t)^2 + \frac{1}{3!} (iL_p t)^3 + \dots \right] f(0)$$

$$e^{iL_p t} f(0) = \left[1 + \dot{p}(0) t \left(\frac{\partial}{\partial p} \right) + \frac{1}{2} (\dot{p}(0) t)^2 \left(\frac{\partial}{\partial p} \right)^2 + \dots \right] f(0)$$

$$f(0 + \dot{p}(0) t) = f(0) + \dot{p}(0) t \left(\frac{\partial f(0)}{\partial p} \right) + \frac{1}{2} (\dot{p}(0) t)^2 \left(\frac{\partial^2 f(0)}{\partial p^2} \right) + \dots$$

the operator iL_p
gives a shift of
the momenta

Hence:

$$e^{iL_p t} f(0) = f(0 + \dot{p}(0) t)$$

The operation iL_r gives a shift of the positions:

$$e^{iL_r t} f(0,0) = f(0, 0 + \dot{r}(0)t)$$

... and the operator iL_p a shift of the momenta:

$$e^{iL_p t} f(0,0) = f(0 + \dot{p}(0)t, 0)$$

This would have been useful if the operators would commute

$$e^{iL t} f(0,0) = e^{(iL_r + iL_p)t} f(0,0) \neq e^{iL_r t} e^{iL_p t} f(0,0)$$

Trotter expansion:

we have the non-commuting operators A and B:

then the following expansion holds:

$$e^{A+B} \neq e^A e^B$$

$$e^{A+B} = \lim_{P \rightarrow \infty} \left(e^{\frac{A}{2P}} e^{\frac{B}{P}} e^{\frac{A}{2P}} \right)^P$$

$$e^{iL_r t} f(0,0) = f(0, 0 + \dot{r}(0)t)$$

$$e^{iL_p t} f(0,0) = f(0 + \dot{p}(0)t, 0)$$

We can apply the Trotter expansion:

$$e^{A+B} = \lim_{P \rightarrow \infty} \left(e^{\frac{A}{2P}} e^{\frac{B}{P}} e^{\frac{A}{2P}} \right)^P$$

$$\Delta t = \frac{t}{P} \quad \frac{iL_r t}{P} = iL_r \Delta t \quad \frac{iL_p t}{2P} = iL_p \frac{\Delta t}{2}$$

These give as operations:

$$e^{iL_r \Delta t} f(p(t), r(t)) = f(p(t), r(t) + \dot{r}(t) \Delta t)$$

gives us a shift of the position:

$$r(t + \Delta t) \rightarrow r(t) + \dot{r}(t) \Delta t$$

$$e^{iL_p \Delta t/2} f(p(t), r(t)) = f\left(p(t) + \dot{p}(t) \frac{\Delta t}{2}, r(t)\right)$$

gives us a shift of the momenta:

$$p(t + \Delta t) \rightarrow p(t) + \dot{p}(t) \frac{\Delta t}{2}$$

$$e^{iL_r \Delta t} \quad r(t + \Delta t) \rightarrow r(t) + \dot{r}(t) \Delta t$$

$$e^{iL_p \frac{\Delta t}{2}} \quad p\left(t + \frac{\Delta t}{2}\right) \rightarrow p(t) + \dot{p}(t) \frac{\Delta t}{2}$$

We can apply the Trotter expansion to integrate M time steps: $t = M \times \Delta t$

$$f(t) = e^{iLt} f(0) = \left(e^{iL_p \frac{\Delta t}{2}} e^{iL_r \Delta t} e^{iL_p \frac{\Delta t}{2}} \right)^M f(0)$$

These give as operations:

$$e^{iL_p \frac{\Delta t}{2}} \quad p\left(\frac{\Delta t}{2}\right) \rightarrow p(0) + \dot{p}(0) \frac{\Delta t}{2}$$

$$e^{iL_r \Delta t} \quad r(\Delta t) \rightarrow r(0) + \dot{r}\left(\frac{\Delta t}{2}\right) \Delta t$$

$$e^{iL_p \frac{\Delta t}{2}} \quad p(\Delta t) \rightarrow p\left(\frac{\Delta t}{2}\right) + \dot{p}(\Delta t) \frac{\Delta t}{2}$$

which gives after one step

$$p(0) \rightarrow p(0) + \left[f(0) + f(\Delta t) \right] \frac{\Delta t}{2}$$

$$r(0) \rightarrow r(0) + \dot{r}\left(\frac{\Delta t}{2}\right) \Delta t = r(0) + v(0) \Delta t + f(0) \frac{\Delta t^2}{2m}$$

which gives after one step

$$r(0) \rightarrow r(0) + \dot{r}\left(\frac{\Delta t}{2}\right)\Delta t = r(0) + v(0)\Delta t + f(0)\frac{\Delta t^2}{2m}$$

$$p(0) \rightarrow p(0) + [f(0) + f(\Delta t)]\frac{\Delta t}{2}$$

Velocity Verlet algorithm

$$r(t + \Delta t) = r(t) + v(t)\Delta t + f(t)\frac{\Delta t^2}{2m}$$

$$v(t + \Delta t) = v(t) + \frac{\Delta t}{2m} [f(t + \Delta t) + f(t)]$$

Velocity Verlet
algorithm:

$$e^{iL_p \frac{\Delta t}{2}} e^{iL_r \Delta t} e^{iL_p \frac{\Delta t}{2}}$$

$$iL_r \Delta t: \quad r(t + \Delta t) \rightarrow r(t) + v(t) \Delta t$$

$$iL_p \frac{\Delta t}{2}: \quad v\left(t + \frac{\Delta t}{2}\right) \rightarrow v(t) + f(t) \frac{\Delta t}{2}$$

Call force(fx)

Do while (t<tmax)

$$iL_p \frac{\Delta t}{2}: \quad v\left(t + \frac{\Delta t}{2}\right) \rightarrow v(t) + f(t) \frac{\Delta t}{2}$$

$$\mathbf{vx} = \mathbf{vx} + \text{delt}t * \mathbf{fx} / 2$$

$$iL_r \Delta t: \quad r(t + \Delta t) \rightarrow r(t) + v(t) \Delta t$$

$$\mathbf{x} = \mathbf{x} + \text{delt}t * \mathbf{vx}$$

Call force(fx)

$$iL_p \frac{\Delta t}{2}: \quad v(t + \Delta t) \rightarrow v\left(t + \frac{\Delta t}{2}\right) + f(t + \Delta t) \frac{\Delta t}{2}$$

$$\mathbf{vx} = \mathbf{vx} + \text{delt}t * \mathbf{fx} / 2$$

enddo

Liouville formulation

Velocity Verlet algorithm

$$r(t + \Delta t) = r(t) + v(t)\Delta t + f(t)\frac{\Delta t^2}{2m}$$
$$v(t + \Delta t) = v(t) + \frac{\Delta t}{2m} [f(t + \Delta t) + f(t)]$$

Transformations:

$$iL_p \Delta t/2: \quad r(t) \rightarrow r(t)$$
$$v(t) \rightarrow v(t) + f(t)\Delta t/2m$$

$$iL_r \Delta t: \quad r(t + \Delta t) \rightarrow r(t) + v(t)\Delta t$$
$$v(t) \rightarrow v(t)$$

$$J_p = \text{Det} \begin{vmatrix} 1 & 0 \\ \left(\frac{\partial f}{\partial r}\right)\frac{\Delta t}{2m} & 1 \end{vmatrix} = 1$$

$$J_r = \text{Det} \begin{vmatrix} 1 & \Delta t \\ 0 & 1 \end{vmatrix} = 1$$



Three subsequent coordinate transformations in either r or v of which the Jacobian is one: Area preserving

4. Molecular dynamics:

4.5 Multiple time steps

Multiple time steps

What to do with “stiff” potentials?



- Fixed bond-length: constraints (Shake)
- Very small time step

$$iL_r \Delta t : r(t + \Delta t) \rightarrow r(t) + v(t) \Delta t$$

$$iL_p \frac{\Delta t}{2} : v\left(t + \frac{\Delta t}{2}\right) \rightarrow v(t) + f(t) \frac{\Delta t}{2}$$

We can split the force is the stiff part and the more slowly changing rest of the forces: $f(t) = f_{Short}(t) + f_{Long}(t)$

This allows us to split the Liouville operator:

$$iL t = iL_r t + iL_{pShort} t + iL_{pLong}$$

The conventional Trotter expansion:

$$iL t = \left[iL_{pLong} \Delta t / 2 \left[iL_r + iL_{pShort} \right] \Delta t iL_{pLong} \Delta t / 2 \right]^M$$

Now we can make another Trotter expansion: $\delta t = \Delta t / m$

$$\left[iL_r + iL_{pShort} \right] \Delta t = \left[iL_{pShort} \delta t / 2 iL_r \delta t iL_{pShort} \delta t / 2 \right]^m$$

The algorithm to solve the equations of motion

$$f(t) = f_{Short}(t) + f_{Long}(t)$$

$$iL t = \left[iL_{pLong} \Delta t/2 \left[iL_r + iL_{pShort} \right] \Delta t iL_{pLong} \Delta t/2 \right]^M$$

$$\left[iL_r + iL_{pShort} \right] \Delta t = \left[iL_{pShort} \delta t/2 iL_r \delta t iL_{pShort} \delta t/2 \right]^m$$

We now have 3 transformations:

$$iL_{pLong} \frac{\Delta t}{2} : v\left(t + \frac{\Delta t}{2}\right) \rightarrow v(t) + f_{Long}(t) \frac{\Delta t}{2}$$

$$iL_{pShort} \frac{\delta t}{2} : v\left(t + \frac{\delta t}{2}\right) \rightarrow v(t) + f_{Short}(t) \frac{\delta t}{2}$$

$$iL_r \delta t : r(t + \delta t) \rightarrow r(t) + v(t) \delta t$$

The steps are first iL_{pLong} then m times iL_{pShort}/iL_r followed by iL_{pLong} again

$$iL_{pLong} \frac{\Delta t}{2} : v\left(t + \frac{\Delta t}{2}\right) \rightarrow v(t) + f_{Long}(t) \frac{\Delta t}{2}$$

Call force(fx_long, f_short)

vx=vx+delt*fx_long/2

Do ddt=1,n

$$iL_{pShort} \frac{\delta t}{2} : v\left(t + \frac{\delta t}{2}\right) \rightarrow v(t) + f_{Short}(t) \frac{\delta t}{2}$$

vx=vx+ddelt*fx_short/2

$$iL_r \delta t : r(t + \delta t) \rightarrow r(t) + v(t) \delta t$$

x=x+ddelt*vx

Call force_short(fx_short)

$$iL_{pShort} \frac{\delta t}{2} : v\left(t + \frac{\delta t}{2}\right) \rightarrow v(t) + f_{Short}(t) \frac{\delta t}{2}$$

vx=vx+ddelt*fx_short/2

enddo

Algorithm 29 (Multiple Time Step)

```
subroutine
+   multi(f_long, f_short)

vx=vx+0.5*delt*f_long
do it=1,n
    vx=vx+0.5*(delt/n)*f_short
    x=x+(delt/n)*vx
    call force_short(f_short)
    vx=vx+0.5*(delt/n)*f_short
enddo
call force_all(f_long, f_short)
vx=vx+0.5*delt*f_long
return
end
```

Multiple time step, `f_long` is
the long-range part and `f_short`
the short-range part of the force
velocity Verlet with time step Δt
loop for the small time step
velocity Verlet with timestep $\Delta t/n$

short-range forces

all forces